

SIX-DIMENSIONAL PAINLEVÉ SYSTEMS AND THEIR PARTICULAR SOLUTIONS IN TERMS OF HYPERGEOMETRIC FUNCTIONS

TAKAO SUZUKI

ABSTRACT. In this article, we propose a class of six-dimensional Painlevé systems which is given as the monodromy preserving deformations of the Fuchsian systems. We also give their particular solutions in terms of hypergeometric functions.

Key Words: Painlevé system, Hypergeometric function, Monodromy.

2000 Mathematics Subject Classification: 34M55, 33C70, 34M35.

1. INTRODUCTION

Recently, higher order generalizations of the sixth Painlevé equation (P_{VI}) has been studied from a viewpoint of the monodromy preserving deformations of Fuchsian systems. It is shown in [10, 11] that any irreducible Fuchsian system can be reduced to finite types of systems by using Katz's two operations, addition and middle convolution [8]. It is also shown in [4] that the isomonodromy deformation equation is invariant under Katz's two operations. Those fact allows us to construct a classification theory of isomonodromy deformation equations.

The Fuchsian systems with two accessory parameters were classified by Kostov [10]. According to it, they are reduced to the systems with the following spectral types:

4 singularities	11, 11, 11, 11
3 singularities	111, 111, 111 22, 1111, 1111 33, 222, 111111

The system with the spectral type $\{11, 11, 11, 11\}$ gives P_{VI} as the monodromy preserving deformation. The other three systems have no deformation parameters, thus we can not derive isomonodromy deformation equations from them.

In general, Fuchsian systems can be classified with the aid of algorithm proposed by Oshima [11]. The systems with four accessory parameters are reduced as follows:

5 singularities	11, 11, 11, 11, 11
4 singularities	21, 21, 111, 111 31, 22, 22, 1111 22, 22, 22, 211

In addition to them, there exist nine systems which have three singularities; we do not list here. Sakai investigated their monodromy preserving deformations systematically and derived the four-dimensional Painlevé systems as follows:

Spectral type	Painlevé system	Ref.
11, 11, 11, 11, 11	Garnier	[3]
21, 21, 111, 111	Fuji-Suzuki	[12, 15]
22, 22, 22, 211	Matrix type	[12]
31, 22, 22, 1111	Sasano	[12]

An aim of this article is to investigate the monodromy preserving deformations of the Fuchsian systems with six accessory parameters. They are reduced as follows:

6 singularities	11, 11, 11, 11, 11, 11		
5 singularities	21, 21, 21, 21, 111	31, 31, 22, 22, 22	
4 singularities	21, 111, 111, 111	22, 22, 211, 211	22, 22, 22, 1111
	31, 22, 211, 1111	31, 31, 1111, 1111	33, 33, 33, 321
	42, 33, 33, 222	51, 33, 222, 222	51, 33, 33, 111111

In addition to them, there exist 24 systems which have three singularities; we do not list here. Among those 12 systems, the following ones have been already investigated:

Spectral type	Painlevé system	Ref.
11, 11, 11, 11, 11, 11	Garnier	[3]
31, 31, 1111, 1111	Fuji-Suzuki-Tsuda	[15]
33, 33, 33, 321	Matrix type	[9]
51, 33, 33, 111111	Sasano	[1]

In this article, we investigate for the other eight Fuchsian systems and derive six-dimensional Painlevé systems.

The other aim of this article is to give particular solutions of the six-dimensional Painlevé systems in terms of hypergeometric functions. It is known that P_{VI} and the four-dimensional Painlevé systems admit particular solutions as follows:

Painlevé system	Rigid system	HGF	Ref.
11, 11, 11, 11	11, 11, 11, 11	${}_2F_1$	[5]
11, 11, 11, 11, 11	21, 21, 21, 21	$P_3(F_1)$	[5]
21, 21, 111, 111	21, 111, 111	${}_3F_2$	[14, 16]

And, for the six-dimensional ones, we obtain the following results:

Painlevé system	Rigid system	HGF	Ref.
11, 11, 11, 11, 11, 11	31, 31, 31, 31, 31	$P_4(F_D)$	[5]
21, 21, 21, 21, 111	31, 31, 22, 211		Sec. 5.1
31, 31, 22, 22, 22	31, 22, 22, 22		Sec. 5.2
21, 111, 111, 111	211, 211, 211		Sec. 5.3
31, 22, 211, 1111	22, 211, 1111	EO_4	Sec. 5.4
31, 31, 1111, 1111	31, 1111, 1111	${}_4F_3$	[14, 16]

This article is organized as follows. In Section 2 and Appendix A, we give explicit formulas of the six-dimensional Painlevé systems. They are expressed as sixth order Hamiltonian systems. In Section 3, we recall the Schlesinger system and its Poisson structure. In Section 4, we discuss derivations of the Painlevé systems from the Schlesinger systems. In Section 5, we give particular solutions of the six-dimensional Painlevé systems in terms of hypergeometric functions. In Appendix B, we recall the four-dimensional Painlevé systems which have been classified by Sakai.

2. LIST OF HAMILTONIANS OBTAINED IN THIS ARTICLE

It is known that P_{VI} can be expressed as the Hamiltonian system

$$t(t-1)\frac{dq}{dt} = \{H_{VI}, q\}, \quad t(t-1)\frac{dp}{dt} = \{H_{VI}, p\},$$

with

$$\begin{aligned} H_{VI} &= H_{VI}(\alpha_0, \alpha_1, \alpha_3, \alpha_4; q, p; t) \\ &= q(q-1)(q-t)p \left(p - \frac{\alpha_1-1}{q-t} - \frac{\alpha_3}{q-1} - \frac{\alpha_4}{q} \right) + \alpha_2(\alpha_0 + \alpha_2)q, \end{aligned}$$

where $\alpha_0 + \alpha_1 + 2\alpha_2 + \alpha_3 + \alpha_4 = 1$. Note that P_{VI} is derived from the Fuchsian system with four singularities and two accessory parameters.

Such a property holds even in higher dimensional cases. The $2n$ -dimensional Painlevé systems which have been derived until now can be expressed as the $2n$ -th order Hamiltonian systems

$$\mathcal{H}^{\mathbf{m}} : \quad t_i(t_i-1)\frac{\partial q_j}{\partial t_i} = \{H_i^{\mathbf{m}}, q_i\}, \quad t_i(t_i-1)\frac{\partial p_j}{\partial t_i} = \{H_i^{\mathbf{m}}, p_j\} \quad (i = 1, \dots, N; j = 1, \dots, n).$$

Here the symbol $\mathbf{m}$ stands for each spectral type. In the case $N = 1$, we denote t_1 and H_1 by t and H respectively. Note that corresponding Fuchsian systems have $N + 3$ singularities and $2n$ accessory parameters.

In this section, we give explicit formulas of the following Hamiltonians:

$$\begin{aligned} &H_i^{21,21,21,21,111}, \quad H_i^{31,31,22,22,22} \quad (i = 1, 2), \\ &H^{21,111,111,111}, \quad H^{31,22,211,1111}, \quad H^{22,22,211,211}, \quad H^{22,22,22,1111}, \quad H^{42,33,33,222}, \quad H^{51,33,222,222}. \end{aligned}$$

The following ones are given in Appendix A:

$$H_i^{11,11,11,11,11,11}, \quad (i = 1, 2, 3), \quad H^{31,31,1111,1111}, \quad H^{33,33,33,321}, \quad H^{51,33,33,111111}.$$

And the following ones are given in Appendix B:

$$H_i^{11,11,11,11,11}, \quad (i = 1, 2), \quad H^{21,21,111,111}, \quad H^{22,22,22,211}, \quad H^{31,22,22,1111}.$$

2.1. Spectral type 21, 21, 21, 21, 111.

$$\begin{aligned} &H_1^{21,21,21,21,111} \\ &= H_1^{11,11,11,11,11,11} + \frac{t_1-1}{t_1-t_2}q_3(q_3-t_2)(q_3-t_1)p_3 \left(p_3 - \frac{\alpha_0+\alpha_3+\alpha_5}{q_3-t_1} - \frac{\alpha_0+\alpha_4-\rho_3}{q_3-t_2} - \frac{\alpha_1}{q_3} \right) \\ &\quad - (\alpha_0 + \alpha_5 + 1)q_3p_3 + \frac{t_1-1}{t_1-t_2}\alpha_2\rho_3q_3 + \frac{1}{t_2}q_2q_3(q_1p_1 + q_2p_2 + \alpha_0 + \alpha_5 + 1)(q_3p_3 + \rho_3) \\ &\quad + q_1q_3p_3(q_1p_1 + q_2p_2 + \alpha_0 + \alpha_5 + 1) + \frac{1}{t_2(t_1-t_2)}q_2q_3\{t_1(t_2-1)p_1 - (t_1-1)t_2p_2\}(q_3p_3 + \rho_3) \\ &\quad + \frac{1}{t_1-t_2}q_3p_3\{-(t_1^2+t_1-2t_2)q_1p_1 - t_1(t_1-1)q_2p_1 + (t_1-1)t_2q_1p_2 + t_1(t_2-1)q_2p_2\} \\ &\quad + \frac{t_1}{t_1-t_2}p_3\{(t_1-1)t_2q_1p_1 - (t_1-1)t_2q_1p_2 + (t_1-t_2)q_3p_1\}, \end{aligned}$$

and

$$\begin{aligned}
H_2^{21,21,21,21,111} &= H_2^{11,11,11,11,11} - \rho_3 q_2 \{(q_2 - 1)p_2 + q_1 p_1 + \alpha_0 + \alpha_5 + 1\} \\
&+ \frac{t_2 - 1}{t_2 - t_1} q_3 (q_3 - t_1)(q_3 - t_2) p_3 \left(p_3 - \frac{\alpha_0 + \alpha_4 - \rho_3 - 1}{q_3 - t_2} - \frac{\alpha_0 + \alpha_3 + \alpha_5 + 1}{q_3 - t_1} - \frac{\alpha_1}{q_3} \right) \\
&- (\alpha_0 + \alpha_5 + 1) q_3 p_3 + \frac{t_2 - 1}{t_2 - t_1} \alpha_2 \rho_3 q_3 + q_2 q_3 p_3 (q_2 p_2 + q_1 p_1 + \alpha_0 + \alpha_5 + 1) \\
&- \frac{t_2 - 1}{t_2 - t_1} q_2 q_3 (p_2 - p_1)(q_3 p_3 + \rho_3) + t_2 q_1 p_3 (q_2 p_2 + q_1 p_1 + \alpha_0 + \alpha_5 + 1) \\
&+ \frac{1}{t_2 - t_1} q_3 p_3 \{(t_2^2 + t_2 - 2t_1)q_2 p_2 + t_2(t_2 - 1)q_1 p_2 - (t_2 - 1)t_1 q_2 p_1 - t_2(t_1 - 1)q_1 p_1\} \\
&+ \frac{t_2}{t_2 - t_1} p_3 \{-t_2(t_1 - 1)q_1 p_2 + (t_2 - 1)t_1 q_1 p_1 - (t_2 - t_1)q_3 p_2\},
\end{aligned}$$

where

$$\alpha_0 = \rho_2, \quad \alpha_1 = \theta_1, \quad \alpha_2 = \theta_2 + \rho_3, \quad \alpha_3 = \theta_4, \quad \alpha_4 = \rho_1 - \rho_2 + 1, \quad \alpha_5 = \theta_3 - 1,$$

and $\theta_1 + \theta_2 + \theta_3 + \theta_4 + \rho_1 + \rho_2 + \rho_3 = 0$.

Remark 2.1. The system $\mathcal{H}^{21,21,21,21,111}$ reduces to $\mathcal{H}^{11,11,11,11,11}$ via a specialization $p_3 = \rho_3 = 0$.

2.2. Spectral type 31, 31, 22, 22, 22.

$$\begin{aligned}
H_i^{31,31,22,22,22} &= H_i^{11,11,11,11,11} - (\alpha_i - \alpha_{i+1}) \frac{t_i}{t_i - t_{i+1}} q_{i+1} \{(t_i - 1)p_i - (t_{i+1} - 1)p_{i+1}\} \\
&- (t_i + 1)q_3^2 p_3^2 + \{\alpha_i + \alpha_3 - 1 + (\alpha_i + \alpha_4)t_i\} q_3 p_3 \\
&+ q_i q_3 p_3 (2q_1 p_1 + 2q_2 p_2 + q_3 p_3 + 2\alpha_0 + \alpha_5 + 1) \\
&- q_3 p_{i+1} (2q_i p_i + q_{i+1} p_{i+1} + 2q_3 p_3 + \alpha_{i+1} + 2\alpha_0 + \alpha_5 + 1) \\
&- 2(t_i + 1)q_i q_3 p_i p_3 + t_i q_{i+1} p_3 (q_3 p_3 - \alpha_i + \alpha_{i+1}) \\
&+ 2t_i q_3 p_i p_3 + \frac{1}{t_i - t_{i+1}} q_3 \{t_i(t_i - 1)p_i^2 - 2t_i(t_{i+1} - 1)p_1 p_2 + (t_i - 1)t_{i+1} p_{i+1}^2\},
\end{aligned}$$

for $i \in \mathbb{Z}/2\mathbb{Z}$, where

$$\alpha_0 = \frac{-\theta_1 + \theta_2 + 2\rho_2}{2}, \quad \alpha_1 = \theta_1, \quad \alpha_2 = \frac{\theta_1 - \theta_2}{2}, \quad \alpha_3 = \theta_4, \quad \alpha_4 = \rho_1 - \rho_2 + 1, \quad \alpha_5 = \theta_3 - 1,$$

and $\theta_1 + \theta_2 + 2\theta_3 + 2\theta_4 + 2\rho_1 + 2\rho_2 = 0$.

Remark 2.2. The system $\mathcal{H}^{31,31,22,22,22}$ reduces to the one $\mathcal{H}^{11,11,11,11,11}$ with $\alpha_1 = \alpha_2$ via a specialization $q_3 = \theta_1 + \theta_2 = 0$.

2.3. Spectral type 21, 111, 111, 111.

$$\begin{aligned}
H^{21,111,111,111} &= H^{21,21,111,111} - \theta_{2,1} t (q_1 - q_2) p_1 + (t - 1) q_3 p_3 - q_2 q_3 p_3 (q_1 p_1 + q_3 p_3 - \alpha_5 + \eta) \\
&+ q_2 q_3 p_1 (q_1 p_1 + q_2 p_2 + q_3 p_3 + \eta) + q_1 q_3 p_3 (2q_1 p_1 + q_2 p_2 + q_3 p_3 + \alpha_1 + \eta) \\
&- q_3 p_1 (q_1 p_1 + q_2 p_2 - \alpha_3 + \eta) - q_3 p_3 (t q_2 p_1 + q_1 p_2) - t p_3 (q_1 - q_2) (q_3 p_3 - \theta_{2,1}),
\end{aligned}$$

where

$$\begin{aligned}\alpha_0 &= -\rho_2, & \alpha_1 &= \theta_{3,2} + \rho_2, & \alpha_2 &= \theta_{3,1} - \theta_{3,2}, & \alpha_3 &= -\theta_{2,1} - \theta_{3,1} - \rho_1, \\ \alpha_4 &= \rho_1 - \rho_3 + 1, & \alpha_5 &= \theta_{2,1} + \rho_3, & \eta &= \theta_{2,2} + \theta_{3,2} + \rho_2 + \rho_3,\end{aligned}$$

and $\theta_1 + \theta_{2,1} + \theta_{2,2} + \theta_{3,1} + \theta_{3,2} + \rho_1 + \rho_2 + \rho_3 = 0$.

Remark 2.3. The system $\mathcal{H}^{21,111,111,111}$ reduces to $\mathcal{H}^{21,21,111,111}$ via a specialization $q_3 = \theta_{2,1} = 0$.

2.4. Spectral type 31, 22, 211, 1111.

$$\begin{aligned}H^{31,22,211,1111} &= H^{21,21,111,111} + \rho_4(\alpha_5 - \eta - \rho_4)q_2 \\ &\quad + H_{VI}(\alpha_5 - \eta - 2\rho_4, \alpha_0, \alpha_2 + \alpha_3 + \alpha_4 + \eta, \alpha_1; q_3, p_3; t) \\ &\quad - \frac{1}{t}q_1q_3(q_3p_3 + \rho_4)(q_3p_3 + \alpha_5 - \eta - \rho_4) \\ &\quad - q_1q_3p_3(2q_1p_1 + q_2p_2 - q_3p_3 + \alpha_1 + \eta) + q_2q_3p_3(2q_1p_1 + q_2p_2 + q_3p_3 + \alpha_5) \\ &\quad + tq_2p_3(q_1p_1 - q_2p_2 - q_3p_3 + \alpha_1) + q_3p_3\{(t+1)q_1p_1 - 2tq_2p_1 + q_1p_2 - q_2p_2\} \\ &\quad - tp_1p_3(q_1 - tq_2),\end{aligned}$$

where

$$\begin{aligned}\alpha_0 &= \theta_{3,2}, & \alpha_1 &= -\theta_2 - \theta_{3,2} - \rho_2 - \rho_4, & \alpha_2 &= \theta_2 + \theta_{3,1} + \rho_2 + \rho_4, \\ \alpha_3 &= -\theta_2 - \theta_{3,1} - \rho_1 - \rho_4, & \alpha_4 &= \rho_1 - \rho_3 + 1, & \alpha_5 &= \theta_2 + \rho_3 + \rho_4, & \eta &= \rho_3 - \rho_4,\end{aligned}$$

and $\theta_1 + 2\theta_2 + \theta_{3,1} + \theta_{3,2} + \rho_1 + \rho_2 + \rho_3 + \rho_4 = 0$.

Remark 2.4. The system $\mathcal{H}^{31,22,211,1111}$ reduces to $\mathcal{H}^{21,21,111,111}$ via a specialization $p_3 = \rho_4 = 0$. It also reduces to $\mathcal{H}^{31,22,22,1111}$ via a specialization $(q_1 - t)p_1 - q_3p_3 - \alpha_2 = \alpha_1 + \alpha_2 = 0$.

2.5. Spectral type 22, 22, 211, 211.

$$\begin{aligned}H^{22,22,211,211} &= \text{tr}\{Q(Q-1)P(Q-t)P - (\alpha_1 - 1)Q(Q-1)P - \alpha_3Q(Q-t)P \\ &\quad - \alpha_4(Q-1)(Q-t)P + \alpha_2(\alpha_0 + \alpha_2)Q + (t-1)R(QP - \alpha_4)\},\end{aligned}$$

with

$$P = \frac{1}{t} \begin{pmatrix} p_1 & -p_2 \\ q_2p_2 + \alpha_2 + \alpha_5 & p_3 \end{pmatrix}, \quad Q = -t \begin{pmatrix} q_1 & 1 \\ -q_2 & q_3 \end{pmatrix}, \quad R = \begin{pmatrix} 0 & -(q_1 - q_3)p_2 - p_1 + p_3 \\ 0 & \alpha_1 + \theta_{3,2} - 1 \end{pmatrix},$$

where

$$\begin{aligned}\alpha_0 &= \theta_2, & \alpha_1 &= -\theta_{3,1} + 1, & \alpha_2 &= -\rho_3, \\ \alpha_3 &= \theta_1 + \theta_2 + \theta_{3,1} + 2\rho_3, & \alpha_4 &= -\theta_1, & \alpha_5 &= -\theta_1 - \theta_2 - \theta_{3,2} - \rho_2,\end{aligned}$$

and $2\theta_1 + 2\theta_2 + \theta_{3,1} + \theta_{3,2} + \rho_1 + \rho_2 + 2\rho_3 = 0$.

Remark 2.5. The system $\mathcal{H}^{22,22,211,211}$ reduces to $\mathcal{H}^{22,22,22,211}$ via a specialization $R = O$.

2.6. Spectral type 22, 22, 22, 1111.

$$H^{22,22,22,1111} = \text{tr}\{Q(Q-1)P(Q-t)P - (\alpha_1-1)Q(Q-1)P - \alpha_3Q(Q-t)P \\ - \alpha_4(Q-1)(Q-t)P + \alpha_2(\alpha_0 + \alpha_2)Q + R(QP + PQ)(Q-t) + (\alpha_0 - \alpha_2 + \rho_4)RQ\},$$

with

$$P = \frac{1}{t} \begin{pmatrix} p_3 & -p_2 \\ q_2 p_2 + \alpha_2 + \alpha_5 & p_1 \end{pmatrix}, \quad Q = -t \begin{pmatrix} q_3 & 1 \\ -q_2 & q_1 \end{pmatrix}, \quad R = \begin{pmatrix} \alpha_2 + \rho_4 & (q_1 - q_3)p_2 + p_1 - p_3 \\ 0 & 0 \end{pmatrix},$$

where

$$\alpha_0 = \theta_2, \quad \alpha_1 = -\theta_3 + 1, \quad \alpha_2 = -\rho_3, \\ \alpha_3 = \theta_1 + \theta_2 + \theta_3 + 2\rho_3, \quad \alpha_4 = -\theta_1, \quad \alpha_5 = -\theta_1 - \theta_2 - \theta_3 - \rho_2,$$

and $2\theta_1 + 2\theta_2 + 2\theta_3 + \rho_1 + \rho_2 + \rho_3 + \rho_4 = 0$.

Remark 2.6. The system $\mathcal{H}^{22,22,22,1111}$ reduces to $\mathcal{H}^{22,22,22,211}$ via a specialization $R = O$.

2.7. Spectral type 42, 33, 33, 222.

$$H^{42,33,33,222}$$

$$= H_{\text{VI}}(-\theta_1 - \theta_2 - \rho_2, -\theta_1 - 2\theta_2 - \theta_3 - \rho_2 - 2\rho_3 + 1, \theta_1 + \theta_3, \theta_1 + \theta_2 + 2\rho_2 + 2\rho_3; q_1, p_1; t) \\ + q_1 p_1 (t p_1 - \rho_2) - \frac{\theta_1 - \theta_2 + 2\rho_2}{2} q_2 p_2 \\ + H_{\text{VI}}\left(\frac{\theta_1 - \theta_2 + 2\rho_2}{2}, \frac{\theta_1 + \theta_2 + 2\theta_3 + 2\rho_2}{2}, -\theta_1 - 2\theta_2 - \theta_3 - 2\rho_3 + 1, -2\rho_2; q_2, p_2; t\right) \\ - 2(t+1)q_3^2 p_3^2 - \{-\theta_1 - \theta_2 - 2\theta_3 + 1 + (3\theta_1 + 3\theta_2 + 2\theta_3 + 4\rho_2 + 4\rho_3 - 1)t\}q_3 p_3 \\ - q_1 q_3 (q_1 p_1 + \theta_2)(q_1 p_1 - \theta_1 - \rho_2) \\ - q_1 q_2 p_1 (q_1 p_1 - \frac{\theta_1 - \theta_2 + 2\rho_2}{2}) + q_2 q_3 p_3 (2q_2 p_2 + q_3 p_3 + \frac{\theta_1 + 3\theta_2 + 2\rho_2 + 4\rho_3}{2}) \\ + t q_3 p_2 (-2q_1 p_1 + 3q_2 p_2 + 4q_3 p_3 + 2\theta_1 + 2\theta_2 + 2\rho_2 + 4\rho_3) \\ + q_2 p_3 (2q_1 p_1 - q_3 p_3 - \frac{\theta_1 - \theta_2 + 2\rho_2}{2}) + 2(t+1)q_3 p_3 (q_1 p_1 - q_2 p_2) \\ - 2t\{(t+1)q_3 p_2^2 + p_1 p_2 (q_1 + q_2) + q_3 p_3 (p_1 - 2p_2)\} + 2t p_1 \{(t+1)p_2 - p_3\},$$

where $3\theta_1 + 3\theta_2 + 2\theta_3 + 2\rho_1 + 2\rho_2 + 2\rho_3 = 0$.

2.8. Spectral type 51, 33, 222, 222.

$$H^{51,33,222,222} = H_{\text{VI}}(\alpha_3, -\alpha_1 - 2\alpha_2 - 2\alpha_3 + 1, \alpha_1, \alpha_3; q_1, p_1; t) \\ + H_{\text{VI}}(\alpha_3, -2\alpha_3 - 2\alpha_4 - \alpha_5 + 1, \alpha_5, \alpha_3; q_2, p_2; t) \\ + H_{\text{VI}}(\alpha_3, -\alpha_0 - 2\alpha_3 - 2\alpha_6 + 1, \alpha_0, \alpha_3; q_3, p_3; t) \\ + \{(q_1 - 1)p_1 + \alpha_2\}\{(q_2 - 1)p_2 + \alpha_4\}(q_1 q_2 + t) \\ + \{(q_2 - 1)p_2 + \alpha_4\}\{(q_3 - 1)p_3 + \alpha_6\}(q_2 q_3 + t) \\ + \{(q_3 - 1)p_3 + \alpha_6\}\{(q_1 - 1)p_1 + \alpha_2\}(q_3 q_1 + t),$$

where

$$\begin{aligned}\alpha_0 &= -\theta_1 - \rho_3 + 1, & \alpha_1 &= -\theta_1 - \theta_{3,2} - \rho_2, & \alpha_2 &= \theta_1 + \theta_{3,1} + \rho_2, \\ \alpha_3 &= \theta_{3,2} + \rho_3, & \alpha_4 &= \frac{\theta_1 + \theta_2}{2}, & \alpha_5 &= \rho_2, & \alpha_6 &= \theta_1 + \rho_1,\end{aligned}$$

and $3\theta_1 + \theta_2 + 2\theta_{3,1} + 2\theta_{3,2} + 2\rho_1 + 2\rho_2 + 2\rho_3 = 0$.

Remark 2.7 ([2]). *This system is also derived from the Drinfeld-Sokolov hierarchy of type $E_6^{(1)}$ by a similarity reduction.*

3. SCHLESINGER SYSTEM

We recall the Schlesinger system and its Poisson structure following [7, 12]. Let $A_1, \dots, A_{N+2} \in M_L(\mathbb{C})$. Consider a system of linear differential equations on $\mathbb{P}^1(\mathbb{C})$

$$\frac{\partial}{\partial x} Y(x) = \sum_{i=1}^{N+2} \frac{A_i}{x - t_i} Y(x), \quad (3.1)$$

with regular singularities $x = t_1, \dots, t_{N+2}, \infty$. We assume that each residue matrix A_i can be diagonalized. The monodromy preserving deformation of the system (3.1) is described as a system of partial differential equations

$$\frac{\partial A_j}{\partial t_i} = \frac{[A_i, A_j]}{t_i - t_j} \quad (j \neq i), \quad \frac{\partial A_i}{\partial t_i} = - \sum_{j=1; j \neq i}^{N+2} \frac{[A_i, A_j]}{t_i - t_j}, \quad (3.2)$$

which is called *the Schlesinger system*. Note that the residue matrix $A_\infty := - \sum_{i=1}^{N+2} A_i$ at $x = \infty$ is a constant matrix.

The system (3.2) can be expressed as a Hamiltonian system

$$\frac{\partial A_j}{\partial t_i} = \{K_i, A_j\}, \quad K_i = \sum_{j=1; j \neq i}^{N+2} \frac{\text{tr} A_i A_j}{t_i - t_j}, \quad (3.3)$$

with the Poisson bracket defined by

$$\{(A_i)_{k,l}, (A_j)_{r,s}\} = \delta_{i,j} \{\delta_{r,l}(A_i)_{k,s} - \delta_{k,s}(A_i)_{r,l}\}, \quad (3.4)$$

where $\delta_{i,j}$ stands for the Kronecker delta. In order to derive the canonical Hamiltonian system from (3.3), we use the method established in [6]. Consider a decomposition of matrices A_i as

$$A_i = B_i C_i, \quad B_i = (b_{k,l}^{(i)})_{k,l} \in M_{L, \text{rank} A_i}(\mathbb{C}), \quad C_i = (c_{l,k}^{(i)})_{l,k} \in M_{\text{rank} A_i, L}(\mathbb{C}).$$

Then we can regard $b_{k,l}^{(i)}$ and $c_{l,k}^{(i)}$ as canonical variables. In fact, the Poisson bracket

$$\{b_{k,l}^{(i)}, c_{l,k}^{(i)}\} = 1, \quad \{\text{otherwise}\} = 0,$$

implies the above one (3.4). Here the symplectic form is expressed as

$$\omega = \sum_{i=1}^{N+2} \text{tr}(dB_i \wedge dC_i) = \sum_{i=1}^{N+2} \sum_{k=1}^L \sum_{l=1}^{\text{rank} A_i} db_{k,l}^{(i)} \wedge dc_{l,k}^{(i)}.$$

Note that the number of accessory parameters of the system (3.1) is generally less than the dimension of a space of matrices $(B_1, C_1, \dots, B_{N+2}, C_{N+2})$. We denote the multiplicity data of eigenvalues of $A_1, \dots, A_{N+2}, A_\infty$, called a *spectral type*, by

$$\{m_{1,1} \dots m_{1,l_1}, \dots, m_{N+2,1} \dots m_{N+2,l_{N+2}}, m_{\infty,1} \dots m_{\infty,l_\infty}\}.$$

Then the number of accessory parameters of (3.1) is given by

$$(N+1)L^2 - \sum_{i=1}^{N+2} \sum_{j=1}^{l_i} m_{i,j}^2 - \sum_{j=1}^{l_\infty} m_{\infty,j}^2 + 2.$$

The following lemma plays an important role in derivation of the Painlevé systems.

Lemma 3.1 ([12]). *Let $G \in GL_L(\mathbb{C})$, $\tilde{B}_i = G^{-1}B_i$ and $\tilde{C}_i = C_iG$. If a condition*

$$\text{tr}\{A_\infty(dG)G^{-1} \wedge (dG)G^{-1}\} = 0,$$

is satisfied, then we obtain

$$\omega = \sum_{i=1}^{N+2} \text{tr}(d\tilde{B}_i \wedge d\tilde{C}_i).$$

Thanks to this lemma, via a gauge transformation $\tilde{A}_i = G^{-1}A_iG$, the system (3.3) with (3.4) is transformed to

$$\mathcal{S}_{L,N} : \quad \frac{\partial \tilde{A}_j}{\partial t_i} = \{K_i, \tilde{A}_j\}, \quad K_i = \sum_{j=1; j \neq i}^{N+2} \frac{\text{tr} \tilde{A}_i \tilde{A}_j}{t_i - t_j},$$

with a symplectic form $\omega = \sum_{i=1}^{N+2} \text{tr}(d\tilde{B}_i \wedge d\tilde{C}_i)$, where

$$\tilde{A}_i = \tilde{B}_i \tilde{C}_i, \quad \tilde{B}_i \in M_{L, \text{rank} \tilde{A}_i}(\mathbb{C}), \quad \tilde{C}_i \in M_{\text{rank} \tilde{A}_i, L}(\mathbb{C}).$$

In the following, we always assume that

- (1) $t_{N+1} = 1$ and $t_{N+2} = 0$;
- (2) $\tilde{A}_{N+2}$ is a upper diagonal matrix;
- (3) $\tilde{A}_\infty := -\sum_{i=1}^{N+2} \tilde{A}_i$ is a lower diagonal matrix.

4. DERIVATION OF THE PAINLEVÉ SYSTEM

In the previous section, we define the Schlesinger system $\mathcal{S}_{L,N}$ as a Hamiltonian system. In this section, we derive six-dimensional Painlevé systems from the systems $\mathcal{S}_{L,N}$ with the following spectral types:

$$\begin{aligned} &\{21, 21, 21, 21, 111\}, \quad \{31, 31, 22, 22, 22\}, \quad \{21, 111, 111, 111\}, \quad \{31, 22, 211, 1111\}, \\ &\{22, 22, 211, 211\}, \quad \{22, 22, 22, 1111\}, \quad \{42, 33, 33, 222\}, \quad \{51, 33, 222, 222\}. \end{aligned}$$

First we discuss a derivation for a spectral type $\{21, 21, 21, 21, 111\}$. The other seven systems can be derived in a similar manner, thus we state only results for them.

4.1. **Spectral type 21, 21, 21, 21, 111.** We consider the system $\mathcal{S}_{3,2}$ with matrices

$$\tilde{A}_i = \begin{pmatrix} 1 \\ b_2^{(i)} \\ b_3^{(i)} \end{pmatrix} \begin{pmatrix} \theta_i - b_2^{(i)} c_2^{(i)} - b_3^{(i)} c_3^{(i)} & c_2^{(i)} & c_3^{(i)} \end{pmatrix} \quad (i = 1, 2, 3),$$

and

$$\tilde{A}_4 = \begin{pmatrix} \theta_4 & a_{12}^{(0)} & a_{13}^{(0)} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \tilde{A}_\infty = \begin{pmatrix} \rho_1 & 0 & 0 \\ a_{21}^{(\infty)} & \rho_2 & 0 \\ a_{31}^{(\infty)} & 0 & \rho_3 \end{pmatrix}.$$

These matrices satisfy a relation

$$\tilde{A}_1 + \tilde{A}_2 + \tilde{A}_3 + \tilde{A}_4 + \tilde{A}_\infty = O. \quad (4.1)$$

Note that a Fuchsian relation $\theta_1 + \theta_2 + \theta_3 + \theta_4 + \rho_1 + \rho_2 + \rho_3 = 0$ is also satisfied. The symplectic form is given by

$$\omega = db_2^{(1)} \wedge dc_2^{(1)} + db_3^{(1)} \wedge dc_3^{(1)} + db_2^{(2)} \wedge dc_2^{(2)} + db_3^{(2)} \wedge dc_3^{(2)} + db_2^{(3)} \wedge dc_2^{(3)} + db_3^{(3)} \wedge dc_3^{(3)}. \quad (4.2)$$

In the following, we derive a six-dimensional canonical coordinate system by substituting the relation (4.1) to the 2-form (4.2).

The (2, 2)-component and the (3, 3)-component of (4.1) are described as

$$b_2^{(1)} c_2^{(1)} + b_2^{(2)} c_2^{(2)} + b_2^{(3)} c_2^{(3)} + \rho_2 = 0, \quad b_3^{(1)} c_3^{(1)} + b_3^{(2)} c_3^{(2)} + b_3^{(3)} c_3^{(3)} + \rho_3 = 0.$$

Substituting them to (4.2), we obtain

$$\omega = d(c_2^{(3)} b_2^{(1)}) \wedge d \frac{c_2^{(1)}}{c_2^{(3)}} + d(c_3^{(3)} b_3^{(1)}) \wedge d \frac{c_3^{(1)}}{c_3^{(3)}} + d(c_2^{(3)} b_2^{(2)}) \wedge d \frac{c_2^{(2)}}{c_2^{(3)}} + d(c_3^{(3)} b_3^{(2)}) \wedge d \frac{c_3^{(2)}}{c_3^{(3)}}. \quad (4.3)$$

Hence we can define a eight-dimensional canonical coordinate system by

$$\begin{aligned} \mu_1 - \mu_3 &= c_2^{(3)} b_2^{(1)}, & \lambda_1 &= \frac{c_2^{(1)}}{c_2^{(3)}}, & \mu_3 &= c_3^{(3)} b_3^{(1)}, & \lambda_3 + \lambda_1 &= \frac{c_3^{(1)}}{c_3^{(3)}}, \\ \mu_2 - \mu_4 &= c_2^{(3)} b_2^{(2)}, & \lambda_2 &= \frac{c_2^{(2)}}{c_2^{(3)}}, & \mu_4 &= c_3^{(3)} b_3^{(2)}, & \lambda_4 + \lambda_2 &= \frac{c_3^{(2)}}{c_3^{(3)}}. \end{aligned}$$

Note that the Hamiltonians K_i ($i = 1, 2$) are expressed as polynomials in the dependent variables μ_j, λ_j ($j = 1, 2, 3, 4$); we do not give their explicit formulas here.

In terms of the variables μ_i, λ_i , we can describe the (3, 2)-component of (4.1) as

$$\mu_3 \lambda_3 + \mu_4 \lambda_4 + \rho_4 = 0.$$

Substituting it to (4.3), we obtain

$$\omega = d\mu_1 \wedge \lambda_1 + d\mu_2 \wedge \lambda_2 + d(\lambda_4 \mu_3) \wedge d \frac{\lambda_3}{\lambda_4}.$$

Hence we can define a six-dimensional canonical coordinate system by

$$p_1 = -\frac{\mu_1}{t_1}, \quad q_1 = -t_1 \lambda_1, \quad p_2 = -\frac{\mu_2}{t_2}, \quad q_2 = -t_2 \lambda_2, \quad p_3 = -\frac{\lambda_4 \mu_3}{t_1}, \quad q_1 = -\frac{t_1 \lambda_3}{\lambda_4}.$$

Let

$$H_1 = K_1 + \frac{q_1 p_1}{t_1} + \frac{q_3 p_3}{t_3}, \quad H_2 = K_2 + \frac{q_2 p_2}{t_2}.$$

Then we arrive at

Theorem 4.1. *The dependent variables p_j, q_j ($j = 1, 2, 3$) defined above are found out to be canonical coordinates of a six-dimensional system with the Poisson bracket*

$$\{p_i, q_j\} = \delta_{i,j}, \quad \{p_i, p_j\} = \{q_i, q_j\} = 0 \quad (i, j = 1, 2, 3).$$

Furthermore, they satisfy the Hamiltonian system $\mathcal{H}^{21,21,21,21,111}$.

In the last, we note that the dependent variables p_j, q_j ($j = 1, 2, 3$) are expressed in terms of components of residue matrices A_i ($i = 1, 2, 3, 4$) as

$$\begin{aligned} t_1 p_1 &= -b_2^{(1)} c_2^{(3)} - b_3^{(1)} c_3^{(3)}, & \frac{q_1}{t_1} &= -\frac{c_2^{(1)}}{c_2^{(3)}}, \\ t_2 p_2 &= -b_2^{(2)} c_2^{(3)} - b_3^{(2)} c_3^{(3)}, & \frac{q_2}{t_2} &= -\frac{c_2^{(2)}}{c_2^{(3)}}, \\ t_1 p_3 &= -\frac{b_3^{(1)}(c_3^{(2)} c_2^{(3)} - c_2^{(2)} c_3^{(3)})}{c_2^{(3)}}, & \frac{q_3}{t_1} &= -\frac{c_3^{(1)} c_2^{(3)} - c_2^{(1)} c_3^{(3)}}{c_3^{(2)} c_2^{(3)} - c_2^{(2)} c_3^{(3)}}. \end{aligned}$$

4.2. Spectral type 31, 31, 22, 22, 22. For this case, we consider the system $\mathcal{S}_{4,2}$ with matrices

$$\begin{aligned} \tilde{A}_i &= \begin{pmatrix} 1 \\ b_2^{(i)} \\ b_3^{(i)} \\ b_4^{(i)} \end{pmatrix} \left(\theta_i - b_2^{(i)} c_2^{(i)} - b_3^{(i)} c_3^{(i)} - b_4^{(i)} c_4^{(i)} \quad c_2^{(i)} \quad c_3^{(i)} \quad c_4^{(i)} \right) \quad (i = 1, 2), \\ \tilde{A}_3 &= \begin{pmatrix} 1 \\ B^{(3)} \end{pmatrix} \left(\theta_3 - C^{(3)} B^{(3)} \quad C^{(3)} \right), \quad B^{(3)} = \begin{pmatrix} b_{31}^{(3)} & b_{32}^{(3)} \\ b_{41}^{(3)} & b_{42}^{(3)} \end{pmatrix}, \quad C^{(3)} = \begin{pmatrix} c_{13}^{(3)} & c_{14}^{(3)} \\ c_{23}^{(3)} & c_{24}^{(3)} \end{pmatrix}, \end{aligned}$$

and

$$\tilde{A}_4 = \begin{pmatrix} \theta_4 & 0 & a_{13}^{(0)} & a_{14}^{(0)} \\ 0 & \theta_4 & a_{23}^{(0)} & a_{24}^{(0)} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad \tilde{A}_\infty = \begin{pmatrix} \rho_1 & 0 & 0 & 0 \\ 0 & \rho_1 & 0 & 0 \\ a_{31}^{(\infty)} & a_{32}^{(\infty)} & \rho_2 & 0 \\ a_{41}^{(\infty)} & a_{42}^{(\infty)} & 0 & \rho_2 \end{pmatrix}.$$

Note that a Fuchsian relation $\theta_1 + \theta_2 + 2\theta_3 + 2\theta_4 + 2\rho_1 + 2\rho_2 = 0$ is satisfied.

Let

$$\begin{aligned}
t_1 p_1 &= \frac{(b_3^{(1)} c_3^{(2)} + b_4^{(1)} c_4^{(2)})(c_{13}^{(3)} c_{24}^{(3)} - c_{14}^{(3)} c_{23}^{(3)})}{(c_3^{(2)} c_{24}^{(3)} - c_4^{(2)} c_{23}^{(3)}) - b_2^{(1)}(c_3^{(2)} c_{14}^{(3)} - c_4^{(2)} c_{13}^{(3)})}, \\
\frac{q_1}{t_1} &= \frac{(c_3^{(1)} c_{24}^{(3)} - c_4^{(1)} c_{23}^{(3)}) - b_2^{(1)}(c_3^{(1)} c_{14}^{(3)} - c_4^{(1)} c_{13}^{(3)})}{c_{13}^{(3)} c_{24}^{(3)} - c_{14}^{(3)} c_{23}^{(3)}}, \\
\frac{q_2}{t_2} &= \frac{(c_3^{(2)} c_{24}^{(3)} - c_4^{(2)} c_{23}^{(3)}) - b_2^{(2)}(c_3^{(2)} c_{14}^{(3)} - c_4^{(2)} c_{13}^{(3)})}{c_{13}^{(3)} c_{24}^{(3)} - c_{14}^{(3)} c_{23}^{(3)}}, \\
t_1 t_2 p_3 &= -\frac{(c_{13}^{(3)} c_{24}^{(3)} - c_{14}^{(3)} c_{23}^{(3)})\{b_3^{(1)}(c_{23}^{(3)} - b_2^{(1)} c_{13}^{(3)}) + b_4^{(1)}(c_{24}^{(3)} - b_2^{(1)} c_{14}^{(3)})\}}{(b_2^{(1)} - b_2^{(2)})\{(c_3^{(2)} c_{24}^{(3)} - c_4^{(2)} c_{23}^{(3)}) - b_2^{(1)}(c_3^{(2)} c_{14}^{(3)} - c_4^{(2)} c_{13}^{(3)})\}}, \\
\frac{q_3}{t_1 t_2} &= -\frac{(b_2^{(1)} - b_2^{(2)})(c_3^{(1)} c_4^{(2)} - c_4^{(1)} c_3^{(2)})}{c_{13}^{(3)} c_{24}^{(3)} - c_{14}^{(3)} c_{23}^{(3)}},
\end{aligned}$$

and

$$t_2 p_2 - \frac{(\theta_1 - \theta_2) t_2 q_1}{2(q_1 q_2 - q_3)} = \frac{(c_3^{(1)} b_3^{(2)} + c_4^{(1)} b_4^{(2)})(c_{13}^{(3)} c_{24}^{(3)} - c_{14}^{(3)} c_{23}^{(3)})}{(c_3^{(1)} c_{24}^{(3)} - c_4^{(1)} c_{23}^{(3)}) - b_2^{(2)}(c_3^{(1)} c_{14}^{(3)} - c_4^{(1)} c_{13}^{(3)})}.$$

We also set

$$H_1 = K_1 + \frac{q_1 p_1}{t_1} + \frac{q_3 p_3}{t_3}, \quad H_2 = K_2 + \frac{q_2 p_2}{t_2} + \frac{q_3 p_3}{t_3}.$$

Then we arrive at

Theorem 4.2. *The dependent variables p_j, q_j ($j = 1, 2, 3$) satisfy the system $\mathcal{H}^{31,31,22,22,22}$.*

4.3. Spectral type 21, 111, 111, 111. For this case, we consider the system $\mathcal{S}_{3,1}$ with matrices

$$\tilde{A}_1 = \begin{pmatrix} 1 \\ b_2^{(1)} \\ b_3^{(1)} \end{pmatrix} \begin{pmatrix} \theta_1 - b_2^{(1)} c_2^{(1)} - b_3^{(1)} c_3^{(1)} & c_2^{(1)} & c_3^{(1)} \end{pmatrix}, \quad \tilde{A}_2 = \begin{pmatrix} 1 & 1 \\ b_{21}^{(2)} & b_{22}^{(2)} \\ b_{31}^{(2)} & b_{32}^{(2)} \end{pmatrix} \begin{pmatrix} c_{11}^{(2)} & c_{12}^{(2)} & c_{13}^{(2)} \\ c_{21}^{(2)} & c_{22}^{(2)} & c_{23}^{(2)} \end{pmatrix},$$

where

$$\begin{pmatrix} c_{11}^{(2)} & c_{12}^{(2)} & c_{13}^{(2)} \\ c_{21}^{(2)} & c_{22}^{(2)} & c_{23}^{(2)} \end{pmatrix} \begin{pmatrix} 1 & 1 \\ b_{21}^{(2)} & b_{22}^{(2)} \\ b_{31}^{(2)} & b_{32}^{(2)} \end{pmatrix} = \begin{pmatrix} \theta_{2,1} & 0 \\ 0 & \theta_{2,2} \end{pmatrix},$$

and

$$\tilde{A}_3 = \begin{pmatrix} \theta_{3,1} & a_{12}^{(0)} & a_{13}^{(0)} \\ 0 & \theta_{3,2} & a_{23}^{(0)} \\ 0 & 0 & 0 \end{pmatrix}, \quad \tilde{A}_\infty = \begin{pmatrix} \rho_1 & 0 & 0 \\ a_{21}^{(\infty)} & \rho_2 & 0 \\ a_{31}^{(\infty)} & a_{32}^{(\infty)} & \rho_3 \end{pmatrix}.$$

Note that a Fuchsian relation $\theta_1 + \theta_{2,1} + \theta_{2,2} + \theta_{3,1} + \theta_{3,2} + \rho_1 + \rho_2 + \rho_3 = 0$ is satisfied.

Let $t = t_1$ and

$$\begin{aligned} tp_1 &= -\frac{b_2^{(1)} c_{22}^{(2)} (c_{13}^{(2)} + c_{23}^{(2)})}{c_{23}^{(2)}}, & \frac{q_1}{t} &= -\frac{c_2^{(1)} c_{23}^{(2)}}{c_{22}^{(2)} (c_{13}^{(2)} + c_{23}^{(2)})}, \\ tp_2 &= -b_3^{(1)} (c_{13}^{(2)} + c_{23}^{(2)}), & \frac{q_2}{t} &= -\frac{c_3^{(1)}}{c_{13}^{(2)} + c_{23}^{(2)}}, \\ tp_3 &= -\frac{b_{21}^{(2)} c_{22}^{(2)} (c_{13}^{(2)} + c_{23}^{(2)})}{c_{23}^{(2)}}, & \frac{q_3}{t} &= -\frac{c_{12}^{(2)} c_{23}^{(2)} - c_{13}^{(2)} c_{22}^{(2)}}{c_{22}^{(2)} (c_{13}^{(2)} + c_{23}^{(2)})}. \end{aligned}$$

We also set

$$H = K_1 + \frac{q_1 p_1}{t} + \frac{q_2 p_2}{t} + \frac{q_3 p_3}{t}.$$

Then we arrive at

Theorem 4.3. *The dependent variables p_j, q_j ($j = 1, 2, 3$) satisfy the system $\mathcal{H}^{21,111,111,111}$.*

4.4. Spectral type 31, 22, 211, 1111. For this case, we consider the system $\mathcal{S}_{4,1}$ with matrices

$$\begin{aligned} \tilde{A}_1 &= \begin{pmatrix} 1 \\ b_2^{(1)} \\ b_3^{(1)} \\ b_4^{(1)} \end{pmatrix} \begin{pmatrix} \theta_1 - b_2^{(1)} c_2^{(1)} - b_3^{(1)} c_3^{(1)} - b_4^{(1)} c_4^{(1)} & c_2^{(1)} & c_3^{(1)} & c_4^{(1)} \end{pmatrix}, \\ \tilde{A}_2 &= \begin{pmatrix} 1 \\ B^{(2)} \end{pmatrix} \begin{pmatrix} \theta_2 - C^{(2)} B^{(2)} & C^{(2)} \end{pmatrix}, \quad B^{(2)} = \begin{pmatrix} b_{31}^{(2)} & b_{32}^{(2)} \\ b_{41}^{(2)} & b_{42}^{(2)} \end{pmatrix}, \quad C^{(2)} = \begin{pmatrix} c_{13}^{(2)} & c_{14}^{(2)} \\ c_{23}^{(2)} & c_{24}^{(2)} \end{pmatrix}, \end{aligned}$$

and

$$\tilde{A}_3 = \begin{pmatrix} \theta_{3,1} & a_{12}^{(0)} & a_{13}^{(0)} & a_{14}^{(0)} \\ 0 & \theta_{3,2} & a_{23}^{(0)} & a_{24}^{(0)} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad \tilde{A}_\infty = \begin{pmatrix} \rho_1 & 0 & 0 & 0 \\ a_{21}^{(\infty)} & \rho_2 & 0 & 0 \\ a_{31}^{(\infty)} & a_{32}^{(\infty)} & \rho_3 & 0 \\ a_{41}^{(\infty)} & a_{42}^{(\infty)} & 0 & \rho_4 \end{pmatrix}.$$

Note that a Fuchsian relation $\theta_1 + 2\theta_2 + \theta_{3,1} + \theta_{3,2} + \rho_1 + \rho_2 + \rho_3 + \rho_4 = 0$ is satisfied.

Let $t = t_1$ and

$$\begin{aligned} tp_1 &= \frac{b_2^{(1)} c_{13}^{(2)} (b_2^{(1)} c_2^{(1)} + \theta_2 + \theta_{3,2} + \rho_2 + \rho_4) + b_2^{(1)} b_4^{(1)} (c_4^{(1)} c_{13}^{(2)} - c_3^{(1)} c_{14}^{(2)})}{c_{23}^{(2)}}, & \frac{q_1}{t} &= \frac{c_{23}^{(2)}}{b_2^{(1)} c_{13}^{(2)}}, \\ tp_2 &= -b_3^{(1)} c_{13}^{(2)} - b_4^{(1)} c_{14}^{(2)}, & \frac{q_2}{t} &= -\frac{c_3^{(1)}}{c_{13}^{(2)}}, \\ tp_3 &= -\frac{b_4^{(1)} (c_{13}^{(2)} c_{24}^{(2)} - c_{14}^{(2)} c_{23}^{(2)})}{b_2^{(1)} c_{13}^{(2)}}, & \frac{q_3}{t} &= -\frac{b_2^{(1)} (c_4^{(1)} c_{13}^{(2)} - c_3^{(1)} c_{14}^{(2)})}{c_{13}^{(2)} c_{24}^{(2)} - c_{14}^{(2)} c_{23}^{(2)}}. \end{aligned}$$

We also set

$$H = K_1 + \frac{q_1 p_1}{t} + \frac{q_2 p_2}{t} + \frac{q_3 p_3}{t}.$$

Then we arrive at

Theorem 4.4. *The dependent variables p_j, q_j ($j = 1, 2, 3$) satisfy the system $\mathcal{H}^{31,22,211,1111}$.*

4.5. **Spectral type** 22, 22, 211, 211. For this case, we consider the system $\mathcal{S}_{4,1}$ with matrices

$$\widetilde{A}_i = \begin{pmatrix} 1 \\ B^{(i)} \end{pmatrix} (\theta_i - C^{(i)} B^{(i)} \quad C^{(i)}), \quad B^{(i)} = \begin{pmatrix} b_{31}^{(i)} & b_{32}^{(i)} \\ b_{41}^{(i)} & b_{42}^{(i)} \end{pmatrix}, \quad C^{(i)} = \begin{pmatrix} c_{13}^{(i)} & c_{14}^{(i)} \\ c_{23}^{(i)} & c_{24}^{(i)} \end{pmatrix} \quad (i = 1, 2),$$

and

$$\widetilde{A}_3 = \begin{pmatrix} \theta_{3,1} & a_{12}^{(0)} & a_{13}^{(0)} & a_{14}^{(0)} \\ 0 & \theta_{3,2} & a_{23}^{(0)} & a_{24}^{(0)} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad \widetilde{A}_\infty = \begin{pmatrix} \rho_1 & 0 & 0 & 0 \\ a_{21}^{(\infty)} & \rho_2 & 0 & 0 \\ a_{31}^{(\infty)} & a_{32}^{(\infty)} & \rho_3 & 0 \\ a_{41}^{(\infty)} & a_{42}^{(\infty)} & 0 & \rho_3 \end{pmatrix}.$$

Note that a Fuchsian relation $2\theta_1 + 2\theta_2 + \theta_{3,1} + \theta_{3,2} + \rho_1 + \rho_2 + 2\rho_3 = 0$ is satisfied.

Let $t = t_1$ and

$$\begin{aligned} p_1 &= b_{31}^{(1)} c_{13}^{(2)} + b_{41}^{(1)} c_{14}^{(2)}, \quad q_1 = \frac{c_{13}^{(1)} c_{24}^{(2)} - c_{14}^{(1)} c_{23}^{(2)}}{c_{13}^{(2)} c_{24}^{(2)} - c_{14}^{(2)} c_{23}^{(2)}}, \\ p_2 &= -\frac{(b_{32}^{(1)} c_{13}^{(2)} + b_{42}^{(1)} c_{14}^{(2)})(c_{23}^{(2)} c_{14}^{(2)} - c_{24}^{(2)} c_{13}^{(2)})}{c_{13}^{(1)} c_{14}^{(2)} - c_{14}^{(1)} c_{13}^{(2)}}, \quad q_2 = -\frac{(c_{23}^{(1)} c_{24}^{(2)} - c_{24}^{(1)} c_{23}^{(2)})(c_{13}^{(1)} c_{14}^{(2)} - c_{14}^{(1)} c_{13}^{(2)})}{(c_{13}^{(2)} c_{24}^{(2)} - c_{14}^{(2)} c_{23}^{(2)})(c_{23}^{(2)} c_{14}^{(2)} - c_{24}^{(2)} c_{13}^{(2)})}, \\ p_3 &= b_{32}^{(1)} c_{23}^{(2)} + b_{42}^{(1)} c_{24}^{(2)}, \quad q_3 = \frac{c_{23}^{(1)} c_{14}^{(2)} - c_{24}^{(1)} c_{13}^{(2)}}{c_{14}^{(2)} c_{23}^{(2)} - c_{13}^{(2)} c_{24}^{(2)}}. \end{aligned}$$

We also set $H = K_1$. Then we arrive at

Theorem 4.5. *The dependent variables p_j, q_j ($j = 1, 2, 3$) satisfy the system $\mathcal{H}^{22,22,211,211}$.*

4.6. **Spectral type** 22, 22, 22, 1111. For this case, we consider the system $\mathcal{S}_{4,1}$ with matrices

$$\widetilde{A}_i = \begin{pmatrix} 1 \\ B^{(i)} \end{pmatrix} (\theta_i - C^{(i)} B^{(i)} \quad C^{(i)}), \quad B^{(i)} = \begin{pmatrix} b_{31}^{(i)} & b_{32}^{(i)} \\ b_{41}^{(i)} & b_{42}^{(i)} \end{pmatrix}, \quad C^{(i)} = \begin{pmatrix} c_{13}^{(i)} & c_{14}^{(i)} \\ c_{23}^{(i)} & c_{24}^{(i)} \end{pmatrix} \quad (i = 1, 2),$$

and

$$\widetilde{A}_3 = \begin{pmatrix} \theta_3 & 0 & a_{13}^{(0)} & a_{14}^{(0)} \\ 0 & \theta_3 & a_{23}^{(0)} & a_{24}^{(0)} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad \widetilde{A}_\infty = \begin{pmatrix} \rho_1 & 0 & 0 & 0 \\ 0 & \rho_2 & 0 & 0 \\ a_{31}^{(\infty)} & a_{32}^{(\infty)} & \rho_3 & 0 \\ a_{41}^{(\infty)} & a_{42}^{(\infty)} & 0 & \rho_4 \end{pmatrix}.$$

Note that a Fuchsian relation $2\theta_1 + 2\theta_2 + 2\theta_3 + \rho_1 + \rho_2 + \rho_3 + \rho_4 = 0$ is satisfied.

Let $t = t_1$,

$$\begin{aligned} p_1 &= -\frac{b_{32}^{(1)}(c_{13}^{(2)} c_{24}^{(2)} - c_{14}^{(2)} c_{23}^{(2)})}{c_{14}^{(2)}}, \quad q_1 = -\frac{(c_{23}^{(1)} c_{14}^{(2)} - c_{24}^{(1)} c_{13}^{(2)})c_{14}^{(2)} - (c_{13}^{(1)} c_{14}^{(2)} - c_{14}^{(1)} c_{13}^{(2)})c_{24}^{(2)}}{c_{14}^{(2)}(c_{13}^{(2)} c_{24}^{(2)} - c_{14}^{(2)} c_{23}^{(2)})}, \\ p_2 &= -\frac{(c_{13}^{(1)} c_{14}^{(2)} - c_{14}^{(1)} c_{13}^{(2)})(c_{14}^{(2)} c_{24}^{(2)} - c_{24}^{(2)} c_{14}^{(2)})}{c_{14}^{(2)} c_{14}^{(2)}(c_{13}^{(2)} c_{24}^{(2)} - c_{14}^{(2)} c_{23}^{(2)})}, \\ p_3 &= \frac{(b_{41}^{(1)} c_{14}^{(2)} + b_{42}^{(1)} c_{24}^{(2)})c_{14}^{(2)} + (b_{31}^{(1)} c_{14}^{(2)} + b_{32}^{(1)} c_{24}^{(2)})c_{13}^{(2)}}{c_{14}^{(2)}}, \quad q_3 = \frac{c_{14}^{(1)}}{c_{14}^{(2)}}. \end{aligned}$$

and

$$p_2 - \frac{\theta_1 + \theta_2 + \theta_{3,1} + \rho_2 + \rho_3}{q_2} = -\frac{(b_{31}^{(1)}c_{14}^{(2)} + b_{32}^{(1)}c_{24}^{(2)})(c_{13}^{(2)}c_{24}^{(2)} - c_{14}^{(2)}c_{23}^{(2)})}{c_{14}^{(1)}c_{24}^{(2)} - c_{24}^{(1)}c_{14}^{(2)}}.$$

We also set $H = K_1$. Then we arrive at

Theorem 4.6. *The dependent variables p_j, q_j ($j = 1, 2, 3$) satisfy the system $\mathcal{H}^{22,22,22,1111}$.*

4.7. **Spectral type** 42, 33, 33, 222. For this case, we consider the system $\mathcal{S}_{6,1}$ with matrices

$$\tilde{A}_i = \begin{pmatrix} 1 \\ B^{(i)} \end{pmatrix} \begin{pmatrix} \theta_i - C^{(i)}B^{(i)} & C^{(i)} \end{pmatrix}, \quad B^{(i)} = \begin{pmatrix} b_{41}^{(i)} & b_{42}^{(i)} & b_{43}^{(i)} \\ b_{51}^{(i)} & b_{52}^{(i)} & b_{53}^{(i)} \\ b_{61}^{(i)} & b_{62}^{(i)} & b_{63}^{(i)} \end{pmatrix}, \quad C^{(i)} = \begin{pmatrix} c_{14}^{(i)} & c_{15}^{(i)} & c_{16}^{(i)} \\ c_{24}^{(i)} & c_{25}^{(i)} & c_{26}^{(i)} \\ c_{34}^{(i)} & c_{35}^{(i)} & c_{36}^{(i)} \end{pmatrix} \quad (i = 1, 2),$$

and

$$\tilde{A}_3 = \begin{pmatrix} \theta_3 & 0 & a_{13}^{(0)} & a_{14}^{(0)} & a_{15}^{(0)} & a_{16}^{(0)} \\ 0 & \theta_3 & a_{23}^{(0)} & a_{24}^{(0)} & a_{25}^{(0)} & a_{26}^{(0)} \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad \tilde{A}_\infty = \begin{pmatrix} \rho_1 & 0 & 0 & 0 & 0 & 0 \\ 0 & \rho_1 & 0 & 0 & 0 & 0 \\ a_{31}^{(\infty)} & a_{32}^{(\infty)} & \rho_2 & 0 & 0 & 0 \\ a_{41}^{(\infty)} & a_{42}^{(\infty)} & 0 & \rho_2 & 0 & 0 \\ a_{51}^{(\infty)} & a_{52}^{(\infty)} & 0 & 0 & \rho_3 & 0 \\ a_{61}^{(\infty)} & a_{62}^{(\infty)} & 0 & 0 & 0 & \rho_3 \end{pmatrix}.$$

Note that a Fuchsian relation $3\theta_1 + 3\theta_2 + 2\theta_3 + 2\rho_1 + 2\rho_2 + 2\rho_3 = 0$ is satisfied.

Let $t = t_1$,

$$\begin{aligned} q_1 &= \frac{(b_{41}^{(1)}b_{42}^{(2)} - b_{42}^{(1)}b_{41}^{(2)})(c_{15}^{(2)}c_{26}^{(2)} - c_{16}^{(2)}c_{25}^{(2)})}{(b_{43}^{(1)} - b_{43}^{(2)})\{b_{41}^{(1)}(c_{15}^{(1)}c_{36}^{(1)} - c_{16}^{(1)}c_{35}^{(1)}) + b_{42}^{(1)}(c_{25}^{(1)}c_{36}^{(1)} - c_{26}^{(1)}c_{35}^{(1)})\}}, \\ tp_2 &= (c_{15}^{(2)}c_{26}^{(2)} - c_{16}^{(2)}c_{25}^{(2)}) \\ &\quad \times \frac{(b_{41}^{(1)}b_{52}^{(1)} - b_{42}^{(1)}b_{51}^{(1)})(b_{41}^{(1)}c_{15}^{(1)} + b_{42}^{(1)}c_{25}^{(1)}) + (b_{41}^{(1)}b_{62}^{(1)} - b_{42}^{(1)}b_{61}^{(1)})(b_{41}^{(1)}c_{16}^{(1)} + b_{42}^{(1)}c_{26}^{(1)})}{(b_{41}^{(1)}c_{15}^{(1)} + b_{42}^{(1)}b_{25}^{(1)})(b_{41}^{(1)}c_{16}^{(2)} + b_{42}^{(1)}c_{26}^{(2)}) - (b_{41}^{(1)}c_{16}^{(1)} + b_{42}^{(1)}b_{26}^{(1)})(b_{41}^{(1)}c_{15}^{(2)} + b_{42}^{(1)}c_{25}^{(2)})}, \\ \frac{q_2}{t} &= -\frac{c_{15}^{(1)}c_{26}^{(1)} - c_{16}^{(1)}c_{25}^{(1)} - c_{25}^{(1)}c_{16}^{(1)} + c_{26}^{(1)}c_{15}^{(1)}}{c_{15}^{(2)}c_{26}^{(2)} - c_{16}^{(2)}c_{25}^{(2)}}, \\ tp_3 &= -(c_{15}^{(2)}c_{26}^{(2)} - c_{16}^{(2)}c_{25}^{(2)}) \\ &\quad \times \frac{(b_{41}^{(1)}b_{52}^{(1)} - b_{42}^{(1)}b_{51}^{(1)})(b_{41}^{(1)}c_{15}^{(2)} + b_{42}^{(1)}c_{25}^{(2)}) + (b_{41}^{(1)}b_{62}^{(1)} - b_{42}^{(1)}b_{61}^{(1)})(b_{41}^{(1)}c_{16}^{(2)} + b_{42}^{(1)}c_{26}^{(2)})}{(b_{41}^{(1)}c_{15}^{(1)} + b_{42}^{(1)}b_{25}^{(1)})(b_{41}^{(1)}c_{16}^{(2)} + b_{42}^{(1)}c_{26}^{(2)}) - (b_{41}^{(1)}c_{16}^{(1)} + b_{42}^{(1)}b_{26}^{(1)})(b_{41}^{(1)}c_{15}^{(2)} + b_{42}^{(1)}c_{25}^{(2)})}, \\ \frac{q_3}{t} &= -\frac{c_{15}^{(1)}c_{26}^{(1)} - c_{16}^{(1)}c_{25}^{(1)}}{c_{15}^{(2)}c_{26}^{(2)} - c_{16}^{(2)}c_{25}^{(2)}}, \end{aligned}$$

and

$$p_1 = \frac{(b_{43}^{(1)} - b_{43}^{(2)})\{b_{41}^{(1)}(c_{15}^{(1)}c_{36}^{(1)} - c_{16}^{(1)}c_{35}^{(1)}) + b_{42}^{(1)}(c_{25}^{(1)}c_{36}^{(1)} - c_{26}^{(1)}c_{35}^{(1)})\}}{(b_{41}^{(1)}c_{15}^{(1)} + b_{42}^{(1)}b_{25}^{(1)})(b_{41}^{(1)}c_{16}^{(2)} + b_{42}^{(1)}c_{26}^{(2)}) - (b_{41}^{(1)}c_{16}^{(1)} + b_{42}^{(1)}b_{26}^{(1)})(b_{41}^{(1)}c_{15}^{(2)} + b_{42}^{(1)}c_{25}^{(2)})} \\ \times \left[\frac{(b_{43}^{(1)} - b_{43}^{(2)})\{c_{34}^{(1)}(c_{15}^{(2)}c_{26}^{(2)} - c_{16}^{(2)}c_{25}^{(2)}) + c_{35}^{(1)}(c_{16}^{(2)}c_{24}^{(2)} - c_{14}^{(2)}c_{26}^{(2)}) + c_{36}^{(1)}(c_{14}^{(2)}c_{25}^{(2)} - c_{15}^{(2)}c_{24}^{(2)})\}}{c_{15}^{(2)}c_{26}^{(2)} - c_{16}^{(2)}c_{25}^{(2)}} - \theta_3 \right].$$

We also set

$$H = K_1 + \frac{q_2 p_2}{t} + \frac{q_3 p_3}{t}.$$

Then we arrive at

Theorem 4.7. *The dependent variables p_j, q_j ($j = 1, 2, 3$) satisfy the system $\mathcal{H}^{42,33,33,222}$.*

4.8. Spectral type 51, 33, 222, 222. For this case, we consider the system $\mathcal{S}_{6,1}$ with matrices

$$\tilde{A}_1 = \begin{pmatrix} 1 \\ B^{(1)} \end{pmatrix} (\theta_1 - C^{(1)} B^{(1)} \quad C^{(1)}), \quad B^{(1)} = \begin{pmatrix} b_{41}^{(1)} & b_{42}^{(1)} & b_{43}^{(1)} \\ b_{51}^{(1)} & b_{52}^{(1)} & b_{53}^{(1)} \\ b_{61}^{(1)} & b_{62}^{(1)} & b_{63}^{(1)} \end{pmatrix}, \quad C^{(1)} = \begin{pmatrix} c_{14}^{(1)} & c_{15}^{(1)} & c_{16}^{(1)} \\ c_{24}^{(1)} & c_{25}^{(1)} & c_{26}^{(1)} \\ c_{34}^{(1)} & c_{35}^{(1)} & c_{36}^{(1)} \end{pmatrix}, \\ \tilde{A}_2 = \begin{pmatrix} 1 \\ b_2^{(2)} \\ b_3^{(2)} \\ b_4^{(2)} \\ b_5^{(2)} \\ b_6^{(2)} \end{pmatrix} (c_1^{(2)} \quad c_2^{(2)} \quad c_3^{(2)} \quad c_4^{(2)} \quad c_5^{(2)} \quad c_6^{(2)}),$$

where $c_1^{(2)} = \theta_1 - b_2^{(2)}c_2^{(2)} - b_3^{(2)}c_3^{(2)} - b_4^{(2)}c_4^{(2)} - b_5^{(2)}c_5^{(2)} - b_6^{(2)}c_6^{(2)}$ and

$$\tilde{A}_3 = \begin{pmatrix} 0 & 0 & a_{13}^{(0)} & a_{14}^{(0)} & a_{15}^{(0)} & a_{16}^{(0)} \\ 0 & 0 & a_{23}^{(0)} & a_{24}^{(0)} & a_{25}^{(0)} & a_{26}^{(0)} \\ 0 & 0 & \theta_{3,1} & 0 & 0 & 0 \\ 0 & 0 & 0 & \theta_{3,1} & 0 & 0 \\ 0 & 0 & 0 & 0 & \theta_{3,2} & 0 \\ 0 & 0 & 0 & 0 & 0 & \theta_{3,2} \end{pmatrix}, \quad \tilde{A}_\infty = \begin{pmatrix} \rho_1 & 0 & 0 & 0 & 0 & 0 \\ 0 & \rho_1 & 0 & 0 & 0 & 0 \\ a_{31}^{(\infty)} & a_{32}^{(\infty)} & \rho_2 & 0 & 0 & 0 \\ a_{41}^{(\infty)} & a_{42}^{(\infty)} & 0 & \rho_2 & 0 & 0 \\ a_{51}^{(\infty)} & a_{52}^{(\infty)} & 0 & 0 & \rho_3 & 0 \\ a_{61}^{(\infty)} & a_{62}^{(\infty)} & 0 & 0 & 0 & \rho_3 \end{pmatrix}.$$

Note that a Fuchsian relation $3\theta_1 + \theta_2 + 2\theta_{3,1} + 2\theta_{3,2} + 2\rho_1 + 2\rho_2 + 2\rho_3 = 0$ is satisfied.

Let $t = t_1$ and

$$\begin{aligned}
p_1 &= \frac{(c_3^{(2)} + c_4^{(2)} b_{43}^{(1)}) \{c_{35}^{(1)} (b_{41}^{(1)} c_{16}^{(1)} + b_{42}^{(1)} c_{26}^{(1)}) - c_{36}^{(1)} (b_{41}^{(1)} c_{15}^{(1)} + b_{42}^{(1)} c_{25}^{(1)})\}}{c_5^{(2)} (b_{41}^{(1)} c_{16}^{(1)} + b_{42}^{(1)} c_{26}^{(1)}) - c_6^{(2)} (b_{41}^{(1)} c_{15}^{(1)} + b_{42}^{(1)} c_{25}^{(1)})}, & q_1 &= -\frac{c_5^{(2)} b_{53}^{(1)} + c_6^{(2)} b_{63}^{(1)}}{c_3^{(2)} + c_4^{(2)} b_{43}^{(1)}}, \\
p_2 &= \frac{(b_3^{(2)} b_{43}^{(1)} - b_4^{(2)}) \{c_1^{(2)} (c_5^{(2)} c_{16}^{(1)} - c_6^{(2)} c_{15}^{(1)}) + c_2^{(2)} (c_5^{(2)} c_{26}^{(1)} - c_6^{(2)} c_{25}^{(1)})\}}{b_{41}^{(1)} (c_5^{(2)} c_{16}^{(1)} - c_6^{(2)} c_{15}^{(1)}) + b_{42}^{(1)} (c_5^{(2)} c_{26}^{(1)} - c_6^{(2)} c_{25}^{(1)})}, & q_2 &= -\frac{b_{41}^{(1)} + b_2^{(2)} b_{42}^{(1)}}{b_3^{(2)} b_{43}^{(1)} - b_4^{(2)}}, \\
tp_3 &= \frac{\{(c_5^{(2)} b_{52}^{(1)} + c_6^{(2)} b_{62}^{(1)}) b_{41}^{(1)} - (c_5^{(2)} b_{51}^{(1)} + c_6^{(2)} b_{61}^{(1)}) b_{42}^{(1)}\} \{(c_5^{(2)} c_{26}^{(1)} - c_6^{(2)} c_{25}^{(1)}) - (c_5^{(2)} c_{16}^{(1)} - c_6^{(2)} c_{15}^{(1)}) b_2^{(2)}\}}{b_{41}^{(1)} (c_5^{(2)} c_{16}^{(1)} - c_6^{(2)} c_{15}^{(1)}) + b_{42}^{(1)} (c_5^{(2)} c_{26}^{(1)} - c_6^{(2)} c_{25}^{(1)})}, \\
\frac{q_3}{t} &= -\frac{c_1^{(2)} b_{42}^{(1)} - c_2^{(2)} c_{41}^{(1)}}{(c_5^{(2)} b_{52}^{(1)} + c_6^{(2)} b_{62}^{(1)}) b_{41}^{(1)} - (c_5^{(2)} b_{51}^{(1)} + c_6^{(2)} b_{61}^{(1)}) b_{42}^{(1)}}.
\end{aligned}$$

We also set

$$H = K_1 + \frac{q_3 p_3}{t}.$$

Then we arrive at

Theorem 4.8. *The dependent variables p_j, q_j ($j = 1, 2, 3$) satisfy the system $\mathcal{H}^{51,33,222,222}$.*

5. PARTICULAR SOLUTIONS

In this section, we investigate particular solutions of six-dimensional Painlevé systems in terms of hypergeometric functions.

5.1. Spectral type 21, 21, 21, 21, 111. Under the system $\mathcal{H}^{21,21,21,21,111}$, we consider a specialization

$$p_1 - \frac{\alpha_1}{q_1} = p_2 = q_3 = \alpha_0 + \alpha_1 + \alpha_5 + 1 = 0.$$

Also we set

$$\frac{y_1}{y_4} = q_1, \quad \frac{y_2}{y_4} = q_2, \quad \frac{y_3}{y_4} = t_2 q_1 p_3,$$

where the variable y_4 satisfies a Pfaff system

$$t_1(t_1 - 1) \frac{\partial}{\partial t_1} \log y_4 = (\alpha_5 + 1) q_1, \quad t_2(t_2 - 1) \frac{\partial}{\partial t_2} \log y_4 = t_2 q_1 p_3 + (\alpha_5 + \rho_3 + 1) q_2.$$

Then we have

Theorem 5.1. *A vector of variables $\mathbf{y} = [y_1, y_2, y_3, y_4]$ satisfies a rigid system*

$$\frac{\partial \mathbf{y}}{\partial t_i} = \left(\frac{M_t^{(i)}}{t_i - t_{i+1}} + \frac{M_1^{(i)}}{t_i - 1} + \frac{M_0^{(i)}}{t_i} \right) \mathbf{y} \quad (i \in \mathbb{Z}/2\mathbb{Z}), \quad (5.1)$$

with matrices

$$M_t^{(1)} = \begin{pmatrix} -\alpha_2 & \alpha_2 & -\alpha_2\rho_3 & 0 \\ -\alpha_1 & \alpha_1 & -\alpha_1\rho_3 & 0 \\ 1 & -1 & \rho_3 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad M_1^{(1)} = \begin{pmatrix} -\alpha_1 + \alpha_5 + 1 & 0 & 0 & \alpha_5 + 1 \\ 0 & -\alpha_1 & \alpha_1\rho_3 & 0 \\ 0 & 0 & \alpha_5 + 1 & 0 \\ \alpha_1 & 0 & 0 & 0 \end{pmatrix},$$

$$M_0^{(1)} = \begin{pmatrix} \alpha_1 - \alpha_3 + 1 & -\alpha_2 & \alpha_2\rho_3 & -\alpha_5 - 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

and

$$M_t^{(2)} = \begin{pmatrix} -\alpha_2 & \alpha_2 & -\alpha_2\rho_3 & 0 \\ -\alpha_1 & \alpha_1 & -\alpha_1\rho_3 & 0 \\ 1 & -1 & \rho_3 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad M_1^{(2)} = \begin{pmatrix} \alpha_2 & 0 & 0 & 0 \\ 0 & \alpha_2 + \alpha_5 + \rho_3 + 1 & 0 & \alpha_5 + \rho_3 + 1 \\ 0 & 1 & \alpha_2 & 1 \\ 0 & -\alpha_2 & 0 & 0 \end{pmatrix},$$

$$M_0^{(2)} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ \alpha_1 & -\alpha_2 - \alpha_3 - \rho_3 + 1 & 0 & -\alpha_5 - \rho_3 - 1 \\ -1 & 0 & -\alpha_2 - \alpha_3 - \rho_3 + 1 & -1 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

Note that, for each $i \in \mathbb{Z}/2\mathbb{Z}$, the system (5.1) is a Fuchsian one with a spectral type 31, 31, 22, 211.

5.2. Spectral type 31, 31, 22, 22, 22. Under the system $\mathcal{H}^{31,31,22,22,22}$, we consider a specialization

$$q_1 = q_2 = q_3 = \alpha_1 - 2\alpha_2 = 0.$$

Note that such a specialization implies $\tilde{A}_1 = O$. We also set

$$\frac{y_1}{y_4} = p_1, \quad \frac{y_2}{y_4} = p_2, \quad \frac{y_3}{y_4} = \frac{\alpha_1 - 2\alpha_2}{2}p_3 + p_1p_2,$$

where the variable y_4 satisfies a Pfaff system

$$(t_i - 1) \frac{\partial}{\partial t_i} \log y_4 = p_i \quad (i \in \mathbb{Z}/2\mathbb{Z}).$$

Then we have

Theorem 5.2. *A vector of variables $\mathbf{y} = {}^t[y_1, y_2, y_3, y_4]$ satisfies a rigid system*

$$\frac{\partial \mathbf{y}}{\partial t_i} = \left(\frac{M_t^{(i)}}{t_i - t_{i+1}} + \frac{M_1^{(i)}}{t_i - 1} + \frac{M_0^{(i)}}{t_i} \right) \mathbf{y} \quad (i \in \mathbb{Z}/2\mathbb{Z}), \quad (5.2)$$

with matrices

$$M_t^{(1)} = \begin{pmatrix} -\frac{1}{2}(\alpha_1 - 2\alpha_2) & \frac{1}{2}(\alpha_1 - 2\alpha_2) & 0 & 0 \\ \frac{1}{2}(\alpha_1 - 2\alpha_2) & -\frac{1}{2}(\alpha_1 - 2\alpha_2) & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

$$M_1^{(1)} = \begin{pmatrix} 2\alpha_0 + \alpha_1 + \alpha_5 + 1 & 0 & 0 & 1 \\ 0 & \frac{1}{2}(\alpha_1 - 2\alpha_2) & -(\alpha_0 + \alpha_2)(\alpha_0 + \alpha_2 + \alpha_5 + 1) & 0 \\ 0 & 1 & 2\alpha_0 + \alpha_1 + \alpha_5 + 1 - \frac{1}{2}(\alpha_1 - 2\alpha_2) & 0 \\ -\frac{1}{4}(2\alpha_0 + \alpha_1)(2\alpha_0 + \alpha_1 + 2\alpha_5 + 2) & 0 & 0 & 0 \end{pmatrix},$$

$$M_0^{(1)} = \begin{pmatrix} \alpha_3 - 1 & 0 & 0 & 0 \\ -\frac{1}{2}(\alpha_1 - 2\alpha_2) & 0 & (\alpha_0 + \alpha_2)(\alpha_0 + \alpha_2 + \alpha_5 + 1) & 0 \\ 0 & 0 & \alpha_3 - 1 & 0 \\ \frac{1}{4}(2\alpha_0 + \alpha_1)(2\alpha_0 + \alpha_1 + 2\alpha_5 + 2) & 0 & 0 & 0 \end{pmatrix},$$

and

$$M_t^{(2)} = E_{12}M_t^{(1)}E_{12}, \quad M_1^{(2)} = E_{12}M_1^{(1)}E_{12}, \quad M_0^{(2)} = E_{12}M_0^{(1)}E_{12}, \quad E_{12} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Note that, for each $i \in \mathbb{Z}/2\mathbb{Z}$, the system (5.2) is a Fuchsian one with a spectral type 31, 22, 22, 22.

5.3. Spectral type 21, 111, 111, 111. Under the system $\mathcal{H}^{21,111,111,111}$, we consider a specialization

$$p_1 = p_2 = p_3 = \eta = 0.$$

Also we set

$$\frac{y_1}{y_4} = \frac{q_1}{t}, \quad \frac{y_2}{y_4} = \frac{q_2}{t}, \quad \frac{y_3}{y_4} = \frac{q_3}{t},$$

where the variable y_4 satisfies a Pfaff system

$$t(t-1)\frac{d}{dt}\log y_4 = (-\alpha_1 + \eta)q_1 - \alpha_5q_2 - \alpha_3 + \eta.$$

Then we have

Theorem 5.3. A vector of variables $\mathbf{y} = {}^t[y_1, y_2, y_3, y_4]$ satisfies a rigid system

$$\frac{d\mathbf{y}}{dt} = \left(\frac{M_1}{t-1} + \frac{M_0}{t} \right) \mathbf{y}, \tag{5.3}$$

with matrices

$$M_1 = \begin{pmatrix} -\alpha_1 + \eta - \theta_{2,1} & -\alpha_1 + \eta & \theta_{2,1} & -\alpha_1 + \eta \\ -\alpha_5 + \theta_{2,1} & -\alpha_5 & -\theta_{2,1} & -\alpha_5 \\ \alpha_3 - \eta & 0 & -\alpha_3 + \eta & 0 \\ -\alpha_3 + \eta & -\alpha_3 + \eta & 0 & -\alpha_3 + \eta \end{pmatrix},$$

$$M_0 = \begin{pmatrix} -\alpha_2 - \eta & \alpha_1 - \eta & 0 & 0 \\ 0 & -\alpha_0 - \alpha_1 - \alpha_2 & 0 & 0 \\ -\alpha_3 + \eta & 0 & \alpha_3 - \eta & 0 \\ \alpha_3 - \eta & \alpha_3 - \eta & 0 & \alpha_3 - \eta \end{pmatrix}.$$

Note that the system (5.3) is a Fuchsian one with a spectral type 211, 211, 211.

5.4. Spectral type 31, 22, 211, 1111. Under the system $\mathcal{H}^{31,22,211,1111}$, we consider a specialization

$$q_1 - \frac{\alpha_1}{p_1} = q_2 = q_3 = \alpha_1 + \alpha_3 - \eta = 0.$$

Note that such a specialization implies $\tilde{A}_1 = O$. We also set

$$\frac{y_1}{y_4} = p_1, \quad \frac{y_2}{y_4} = p_2, \quad \frac{y_3}{y_4} = tp_1p_3,$$

where the variable y_4 satisfies a Pfaff system

$$t(t-1)\frac{d}{dt}\log y_4 = tp_1 + tp_2 + (\alpha_3 - \rho_4)t + (\alpha_3 + \alpha_4 + \alpha_5)(t-1)$$

Then we have

Theorem 5.4. A vector of variables $\mathbf{y} = {}^t[y_1, y_2, y_3, y_4]$ satisfies a rigid system

$$\frac{d\mathbf{y}}{dt} = \left(\frac{M_1}{t-1} + \frac{M_0}{t} \right) \mathbf{y}, \tag{5.4}$$

with matrices

$$M_1 = \begin{pmatrix} -\alpha_1 - \rho_4 & \alpha_5 & -\rho_4(\alpha_5 - \eta - \rho_4) & 1 \\ -\alpha_1 & \alpha_5 - \rho_4 & 0 & 1 \\ 1 & -1 & \alpha_5 - \eta - \rho_4 & 0 \\ -\alpha_1(\alpha_1 - \eta) & \alpha_1\alpha_5 - (\alpha_5 - \rho_4)(\eta + \rho_4) & -\alpha_1\rho_4(\alpha_5 - \eta - \rho_4) & \alpha_1 - \eta - \rho_4 \end{pmatrix},$$

$$M_0 = \begin{pmatrix} -\alpha_0 & 0 & 0 & 0 \\ \alpha_1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ \alpha_1(\alpha_1 - \eta) & -\alpha_1\alpha_5 + (\alpha_5 - \rho_4)(\eta + \rho_4) & \alpha_1\rho_4(\alpha_5 - \eta - \rho_4) & \alpha_3 + \alpha_4 + \alpha_5 \end{pmatrix}.$$

Note that the system (5.4) is a Fuchsian one with a spectral type 22, 211, 1111.

APPENDIX A. HAMILTONIANS OF THE SIX-DIMENSIONAL PAINLEVÉ SYSTEM

In this section, we recall the Hamiltonian of the six-dimensional Painlevé systems which have been already derived in [3, 15, 9, 1].

A.1. Spectral type 11, 11, 11, 11, 11, 11 (Garnier system).

$$\begin{aligned}
H_i^{11,11,11,11,11,11} = & H_{VI}(\alpha_6 + 1, \alpha_{i+1} + \alpha_{i-1} + \alpha_4, \alpha_5, \alpha_i; q_i, p_i; t_i) \\
& + q_i q_{i+1} p_{i+1} (2q_i p_i + q_{i+1} p_{i+1} + 2\alpha_0 + \alpha_6 + 1) \\
& - \frac{1}{t_i - t_{i+1}} q_i q_{i+1} \{t_i(t_i - 1)p_i^2 - 2t_i(t_{i+1} - 1)p_i p_{i+1} + (t_i - 1)t_{i+1} p_{i+1}^2\} \\
& + \alpha_i \frac{t_i}{t_i - t_{i+1}} q_{i+1} \{(t_i - 1)p_i - (t_{i+1} - 1)p_{i+1}\} - \alpha_{i+1} \frac{(t_i - 1)t_{i+1}}{t_i - t_{i+1}} q_i (p_i - p_{i+1}) \\
& + q_i q_{i-1} p_{i-1} (2q_i p_i + q_{i-1} p_{i-1} + 2\alpha_0 + \alpha_6 + 1) \\
& - \frac{1}{t_i - t_{i-1}} q_i q_{i-1} \{t_i(t_i - 1)p_i^2 - 2t_i(t_{i-1} - 1)p_i p_{i-1} + (t_i - 1)t_{i-1} p_{i-1}^2\} \\
& + \alpha_i \frac{t_i}{t_i - t_{i-1}} q_{i-1} \{(t_i - 1)p_i - (t_{i-1} - 1)p_{i-1}\} - \alpha_{i-1} \frac{(t_i - 1)t_{i-1}}{t_i - t_{i-1}} q_i (p_i - p_{i-1}),
\end{aligned}$$

for $i \in \mathbb{Z}/3\mathbb{Z}$, where $2\alpha_0 + \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6 = 0$.

A.2. Spectral type 31, 31, 1111, 1111 (Fuji-Suzuki-Tsuda system).

$$\begin{aligned}
H^{31,31,1111,1111} = & H_{VI}(-\alpha_1 + \eta, \alpha_2 + \alpha_4 + \alpha_6, \alpha_0, \alpha_3 + \alpha_5 + \alpha_7 - \eta; q_i, p_i; t) \\
& + H_{VI}(-\alpha_3 + \eta, \alpha_4 + \alpha_6, \alpha_0 + \alpha_2, \alpha_1 + \alpha_5 + \alpha_7 - \eta; q_i, p_i; t) \\
& + H_{VI}(-\alpha_5 + \eta, \alpha_6, \alpha_0 + \alpha_2 + \alpha_4, \alpha_1 + \alpha_3 + \alpha_7 - \eta; q_i, p_i; t) \\
& + (q_1 - 1)(q_2 - t) \{(q_1 p_1 + \alpha_1) p_2 + p_1 (p_2 q_2 + \alpha_3)\} \\
& + (q_1 - 1)(q_3 - t) \{(q_1 p_1 + \alpha_1) p_3 + p_1 (p_3 q_3 + \alpha_5)\} \\
& + (q_2 - 1)(q_3 - t) \{(q_2 p_2 + \alpha_3) p_3 + p_2 (p_3 q_3 + \alpha_5)\},
\end{aligned}$$

where $\alpha_0 + \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6 + \alpha_7 = 1$.

A.3. Spectral type 33, 33, 33, 321 (Matrix-type sixth Painlevé system).

$$\begin{aligned}
H^{33,33,33,321} = & \text{tr}\{Q(Q - 1)P(Q - t)P - (\alpha_1 - 1)Q(Q - 1)P \\
& - \alpha_3 Q(Q - t)P - \alpha_4 (Q - 1)(Q - t)P + \alpha_2 (\alpha_0 + \alpha_2)Q\}
\end{aligned}$$

where $\alpha_0 + \alpha_1 + 2\alpha_2 + \alpha_3 + \alpha_4 = 1$ and

$$\begin{aligned}
P = & \frac{1}{t} \begin{pmatrix} \frac{1}{3}p_1 + q_1^2 p_3 & \frac{2}{3}q_2 p_2 + (q_3 - q_1^3)p_3 - 2\alpha_2 - 2\alpha_5 & \frac{1}{3}(q_3 - q_1^3)p_2 + 2q_2^2 p_3 \\ \frac{1}{3}p_2 & \frac{1}{3}p_1 + (q_2 + q_1^2)p_3 & \frac{1}{3}q_2 p_2 + (q_3 - q_1^3)p_3 - \alpha_2 - \alpha_5 \\ p_3 & \frac{1}{3}p_2 & \frac{1}{3}p_1 - (q_2 - q_1^2)p_3 \end{pmatrix}, \\
Q = & -t \begin{pmatrix} q_1 & 2q_2 & q_3 - q_1^3 \\ 1 & q_1 & q_2 \\ 0 & 1 & q_1 \end{pmatrix}.
\end{aligned}$$

A.4. Spectral type 51, 33, 33, 111111 (Sasano system).

$$\begin{aligned}
H^{51,33,33,111111} = & H_{\text{VI}}(\alpha_0, \alpha_1, \alpha_3 + 2\alpha_4 + \alpha_5 + 2\alpha_6 + \alpha_7, \alpha_3 + \alpha_5 + \alpha_8; q_1, p_1; t) \\
& + H_{\text{VI}}(\alpha_0 + 2\alpha_2 + \alpha_3, \alpha_1 + \alpha_3, 2\alpha_4 + \alpha_5 + 2\alpha_6 + \alpha_7, \alpha_5 + \alpha_8; q_2, p_2; t) \\
& + H_{\text{VI}}(\alpha_0 + 2\alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5, \alpha_1 + \alpha_3 + \alpha_5, 2\alpha_6 + \alpha_7, \alpha_8; q_3, p_3; t) \\
& + 2(q_1 - t)p_1q_2\{(q_2 - 1)p_2 + \alpha_4\} + 2(q_1 - t)p_1q_3\{(q_3 - 1)p_3 + \alpha_6\} \\
& + 2(q_2 - t)p_2q_3\{(q_3 - 1)p_3 + \alpha_6\},
\end{aligned}$$

where $\alpha_0 + \alpha_1 + 2\alpha_2 + 2\alpha_3 + 2\alpha_4 + 2\alpha_5 + 2\alpha_6 + \alpha_7 + \alpha_8 = 1$.

APPENDIX B. HAMILTONIANS OF THE FOUR-DIMENSIONAL PAINLEVÉ SYSTEM

In this section, we recall the Hamiltonian of the four-dimensional Painlevé systems which have been classified by Sakai [12].

B.1. Spectral type 11, 11, 11, 11, 11 (Garnier system).

$$\begin{aligned}
H_i^{11,11,11,11,11} = & H_{\text{VI}}(\alpha_5 + 1, \alpha_{i+1} + \alpha_3, \alpha_4, \alpha_i; q_i, p_i; t_i) + q_1q_2p_{i+1}(2q_ip_i + q_{i+1}p_{i+1} + 2\alpha_0 + \alpha_5 + 1) \\
& - \frac{1}{t_i - t_{i+1}}q_1q_2\{t_i(t_i - 1)p_i^2 - 2t_i(t_{i+1} - 1)p_1p_2 + (t_i - 1)t_{i+1}p_{i+1}^2\} \\
& + \alpha_i \frac{t_i}{t_i - t_{i+1}}q_{i+1}\{(t_i - 1)p_i - (t_{i+1} - 1)p_{i+1}\} - \alpha_{i+1} \frac{(t_i - 1)t_{i+1}}{t_i - t_{i+1}}q_i(p_i - p_{i+1}),
\end{aligned}$$

for $i \in \mathbb{Z}/2\mathbb{Z}$, where $2\alpha_0 + \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 = 0$.

B.2. Spectral type 21, 21, 111, 111 (Fuji-Suzuki system).

$$\begin{aligned}
H^{21,21,111,111} = & H_{\text{VI}}(-\alpha_1 + \eta, \alpha_2, \alpha_0 + \alpha_4, \alpha_3 + \alpha_5 - \eta; q_1, p_1; t) \\
& + H_{\text{VI}}(-\alpha_5 + \eta, \alpha_0 + \alpha_2, \alpha_4, \alpha_1 + \alpha_3 - \eta; q_2, p_2; t) \\
& + (q_1 - t)(q_2 - 1)\{(q_1p_1 + \alpha_1)p_2 + p_1(q_2p_2 + \alpha_5)\},
\end{aligned}$$

where $\alpha_0 + \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 = 1$.

B.3. Spectral type 22, 22, 22, 211 (Matrix-type sixth Painlevé system).

$$\begin{aligned}
H^{22,22,22,211} = & \text{tr}\{Q(Q - 1)P(Q - t)P - (\alpha_1 - 1)Q(Q - 1)P \\
& - \alpha_3Q(Q - t)P - \alpha_4(Q - 1)(Q - t)P + \alpha_2(\alpha_0 + \alpha_2)Q\},
\end{aligned}$$

where $\alpha_0 + \alpha_1 + 2\alpha_2 + \alpha_3 + \alpha_4 = 1$ and

$$P = \frac{1}{t} \begin{pmatrix} \frac{1}{2}p_1 & -p_2 \\ q_2p_2 + \alpha_2 + \alpha_5 & \frac{1}{2}p_1 \end{pmatrix}, \quad Q = -t \begin{pmatrix} q_1 & 1 \\ -q_2 & q_1 \end{pmatrix}.$$

B.4. Spectral type 31, 22, 22, 1111 (Sasano system).

$$\begin{aligned}
H^{31,22,22,1111} = & H_{\text{VI}}(\alpha_0, \alpha_1, \alpha_3 + 2\alpha_4 + \alpha_5, \alpha_3 + \alpha_6; q_1, p_1; t) \\
& + H_{\text{VI}}(\alpha_0 + 2\alpha_2 + \alpha_3, \alpha_1 + \alpha_3, \alpha_5, \alpha_6; q_2, p_2; t) \\
& + 2(q_1 - t)p_1q_2\{(q_2 - 1)p_2 + \alpha_4\},
\end{aligned}$$

where $\alpha_0 + \alpha_1 + 2\alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6 = 1$.

ACKNOWLEDGEMENT

The author would like to express his gratitude to Dr. Kenta Fuji, Mr. Keisuke Inoue and Mr. Keisuke Shinomiya for fruitful discussion. The author is also grateful to Professors Kazuki Hiroe, Hiroshi Kawakami, Hajime Nagoya, Masatoshi Noumi, Toshio Oshima, Hidetaka Sakai and Yasuhiko Yamada for valuable discussions and advices.

REFERENCES

- [1] K. Fuji, K. Inoue, K. Shinomiya and T. Suzuki, Higher order Painlevé system of type $D_{2n+2}^{(1)}$ and monodromy preserving deformation, Preprint (arXiv:1011.0276).
- [2] K. Fuji and T. Suzuki, Coupled Painlevé VI system with $E_6^{(1)}$ -symmetry, J. Phys. A: Math. Theor. **42** (2009) 145205.
- [3] R. Garnier, Sur des équations différentielles du troisième ordre dont l'intégrale est uniforme et sur une classe d'équations nouvelles d'ordre supérieur dont l'intégrale générale a ses points critiques fixes, Ann. Sci. École Norm. Sup. **29** (1912), 1-126.
- [4] Y. Haraoka and G. M. Filipuk, Middle convolution and deformation for Fuchsian systems, J. Lond. Math. Soc. **76** (2007) 438-450.
- [5] K. Iwasaki, H. Kimura, S. Shimomura and M. Yoshida, From Gauss to Painlevé — A Modern Theory of Special Functions, Aspects of Mathematics **E16** (Vieweg, 1991).
- [6] M. Jimbo, T. Miwa, Y. Mori and M. Sato, Density matrix of an impenetrable Bose gas and the fifth Painlevé transcendent, Physica **1D** (1980), 80-158.
- [7] M. Jimbo, T. Miwa and K. Ueno, Monodromy preserving deformation of linear ordinary differential equations with rational coefficients I, Physica **2D** (1981), 306-352.
- [8] N. M. Katz, Rigid Local Systems, Annals of Mathematics Studies **139** (Princeton University Press, 1995).
- [9] H. Kawakami, Private communication.
- [10] V. P. Kostov, The Deligne-Simpson problem for zero index of rigidity, Perspective in Complex Analysis, Differential Geometry and Mathematical Physics (World Scientific 2001), 1-35.
- [11] T. Oshima, Classification of Fuchsian systems and their connection problem, preprint (arXiv:0811.2916).
- [12] H. Sakai, Isomonodromic deformation and 4-dimensional Painlevé type equations, UTMS **2010-17** (Univ. of Tokyo 2010) 1-21.
- [13] C. T. Simpson, Products of Matrices, Canadian Math. Soc. Conference Proceedings **12** (AMS 1991), 157-185.
- [14] T. Suzuki, A particular solution of a Painlevé system in terms of the hypergeometric function ${}_{n+1}F_n$, SIGMA **6** (2010) 078.
- [15] T. Tsuda, UC hierarchy and monodromy preserving deformation, J. reine angew. Math., in press.
- [16] T. Tsuda, Hypergeometric solution of a certain polynomial Hamiltonian system of isomonodromy type, Quart. J. Math. **63** (2012) 489-505.

DEPARTMENT OF MATHEMATICS, KINKI UNIVERSITY, 3-4-1, KOWAKAE, HIGASHI-OSAKA, OSAKA 577-8502, JAPAN
E-mail address: suzuki@math.kindai.ac.jp